

ON THE WELL-POSEDNESS OF THE SCHRÖDINGER-KORTEWEG-DE VRIES SYSTEM

ZIHUA GUO, YUZHAO WANG

ABSTRACT. We prove that the Cauchy problem for the Schrödinger-Korteweg-de Vries system is locally well-posed for the initial data belonging to the Sobolev spaces $L^2(\mathbb{R}) \times H^{-3/4}(\mathbb{R})$. The new ingredient is that we use the $\bar{F}^s$ type space, introduced by the first author in [10], to deal with the KdV part of the system and the coupling terms. In order to overcome the difficulty caused by the lack of scaling invariance, we prove uniform estimates for the multiplier. This result improves the previous one by Corcho and Linares [6].

1. INTRODUCTION

In this paper, we consider the Cauchy problem for the Schrödinger-Korteweg-de Vries (NLS-KdV) system

$$\begin{cases} i\partial_t u + \partial_x^2 u = \alpha uv + \beta |u|^2 u, & t, x \in \mathbb{R}, \\ \partial_t v + \partial_x^3 v + \frac{1}{2} \partial_x (v^2) = \gamma \partial_x (|u|^2), \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), \end{cases} \quad (1.1)$$

where $u(x, t)$ is a complex-valued function, $v(x, t)$ is a real-valued function and α, β, γ are real constants, (u_0, v_0) are given initial data belonging to $H^{s_1} \times H^{s_2}$. Our main motivation of this paper is inspired by the work of Corcho and Linares [6], and the work of the first named author [10].

The system (1.1) is an important model in fluid mechanics and plasma physics that governs the interactions between short-wave and long wave. The case $\beta = 0$ appears in the study of resonant interaction between short and long capillary-gravity waves on water of uniform finite depth, in plasma physics and in a diatomic lattice system (See [6] and reference therein for more introduction).

Before we state our main results, we recall first the early results on this system. M. Tsutsumi [21] obtained global well-posedness (GWP) for data $(u_0, v_0) \in H^{s+1/2}(\mathbb{R}) \times H^s(\mathbb{R})$ with $s \in \mathbb{Z}_+$. In the resonant case ($\beta = 0$) Guo and Miao [8] proved GWP in the natural energy space $H^s(\mathbb{R}) \times H^s(\mathbb{R})$ with $s \in \mathbb{Z}_+$. Bekiranov, Ogawa and Ponce [1] proved local well-posedness (LWP) in $H^s(\mathbb{R}) \times H^{s-1/2}(\mathbb{R})$ for $s \geq 0$, and this result was improved to $L^2 \times H^{-3/4+}$ by Corcho and Linares [6] which seems to be sharp except $L^2 \times H^{-3/4}$ in view of the results for the two single equations in (1.1) (See [17], [4], [16]). Pecher [18] obtained GWP in $H^s \times H^s$ for $s > 3/5$ ($\beta = 0$) and $s > 2/3$ ($\beta \neq 0$) by using the ideas of I-method [5]. Some generalized interaction equations were considered in [2]. In this paper, we prove the following results:

2000 *Mathematics Subject Classification.* 35Q55.

Key words and phrases. Local well-posedness, Schrödinger-Korteweg-de Vries system.

Theorem 1.1. *Assume $u_0 \in L^2$, $v_0 \in H^{-3/4}$. Then*

(a) *Existence.* *There exist $T = T(\|u_0\|_{L^2}, \|v_0\|_{H^{-3/4}}) > 0$ and a solution u to the Cauchy problem (1.1) satisfying*

$$u \in X_{\tau=-\xi^2}^{0,1/2+}(T) \subset C([-T, T] : L^2), v \in \bar{F}_{\tau=\xi^3}^{-3/4}(T) \subset C([-T, T] : H^{-3/4}).$$

(b) *Uniqueness.* *The solution mapping $S_T : (u_0, v_0) \rightarrow (u, v)$ is the unique extension of the classical solution $(H^\infty, H^\infty) \rightarrow C([-T, T] : H^\infty \times H^\infty)$.*

(c) *Lipschitz continuity.* *For any $R > 0$, the mapping $(u_0, v_0) \rightarrow (u, v)$ is Lipschitz continuous from $\{(u_0, v_0) \in L^2 \times H^{-3/4} : \|u_0\|_{L^2} + \|v_0\|_{H^{-3/4}} < R\}$ to $C([-T, T] : L^2 \times H^{-3/4})$.*

We describe briefly our ideas in proving Theorem 1.1. We also use the scheme as in [6] which is the same spirit as the one by Ginibre, Y. Tsutsumi and Velo [7] for the Zakharov system. The basic idea is that for the second equation in (1.1) we use the $\bar{F}^s$ space that was used by the first named author [10] for the KdV equation, but there is an essential difficulty. For the KdV equation, one can assume the initial data v_0 has a small norm by using the scaling transform. However, for the NLS-KdV system we don't have such an invariant scaling transform. In order to deal with large initial data, we overcome this difficulty by the following way. We observe that the single nonlinear Schrödinger equation with cubic term ($|u|^2 u$)

$$i\partial_t u + \partial_x^2 u = \beta |u|^2 u$$

is L^2 -subcritical. On the other-hand, we also see from [6] that to control the coupling term uv one need less regularity than $H^{-3/4}$ of v . Then we expect that the first equation can be handled without scaling. Thus we scale the system (1.1) according to the second equation as following: if (u, v) solve the system (1.1) with initial data (u_0, v_0) , then we see

$$u_\lambda(t, x) = \lambda^2 u(\lambda^3 t, \lambda x), \quad \phi_1(x) = \lambda^2 u_0(\lambda x), \quad (1.2)$$

$$v_\lambda(t, x) = \lambda^2 v(\lambda^3 t, \lambda x), \quad \phi_2(x) = \lambda^2 v_0(\lambda x), \quad (1.3)$$

satisfy the following system

$$\begin{cases} i\partial_t u + \lambda \partial_x^2 u = \lambda uv + \lambda^{-1} |u|^2 u, & t, x \in \mathbb{R}, \\ \partial_t v + \partial_x^3 v + \frac{1}{2} \partial_x(v^2) = \partial_x(|u|^2), \\ u(0, x) = \phi_1(x), v(0, x) = \phi_2(x). \end{cases} \quad (1.4)$$

It is easy to see that by taking $0 < \lambda \ll 1$, we have $\|\phi_1\|_{L^2} = \|\lambda^2 u_0(\lambda x)\|_{L^2} = \lambda^{3/2} \|u_0\|_2 \leq R$ and $\|\phi_2\|_{H^{-3/4}} = \|\lambda^2 v_0(\lambda x)\|_{H^{-3/4}} \leq 2\lambda^{3/4} \|v_0\|_{H^{-3/4}} = \epsilon_0 \ll 1$. Therefore, it reduces to study the system (1.4) under condition that $0 < \lambda \leq 1$ and the following condition

$$\|\phi_1\|_{L^2} \leq R, \quad \|\phi_2\|_{H^{-3/4}} = \epsilon_0 \ll 1. \quad (1.5)$$

where ϵ_0 is a absolutely constant will be defined later. We will prove well-posedness for (1.4)-(1.5) in $[0, T]$ for some $T = T(R, \lambda) > 0$. By the scaling we obtain local well-posedness for the original system (1.1).

In our proof the condition $0 < \lambda \leq 1$ in (1.4) is crucial. Heuristically, the propagation speed for the first equation is $\lambda\xi$, and that for the second equation is ξ^2 . Then we see that the two waves u, v has a separate speed in high frequency $|\xi| \gtrsim 1$ uniformly for $0 < \lambda \leq 1$. Thus the resonance and coherence can not be simultaneously large (Also see [6] for the case $\lambda = 1$). This is key to control the

coupled wave interactions. Technically, we will prove uniform estimates for the multiplier associated to the coupled terms for all $0 < \lambda \leq 1$. Our proof for the coupled terms is different from those in [6], but with basically the same ideas. We will use the ideas developed by Tao [19], Ionescu and Kenig [12], and the first-named author [9], but in this paper we need to deal with two different wave forms which has independent interest.

At the end of this section we introduce some notations. In Section 2, we prove some L^2 bilinear estimates which will be used to prove the bilinear estimates for the coupling terms in Section 3. In Section 4, we prove Theorem 1.1.

Notations. Throughout this paper we fix $0 < \lambda \leq 1$. We will use C and c to denote constants which are independent of λ and not necessarily the same at each occurrence. For $x, y \in \mathbb{R}$, $x \sim y$ means that there exist $C_1, C_2 > 0$ such that $C_1|x| \leq |y| \leq C_2|x|$. For $f \in \mathcal{S}'$ we denote by $\widehat{f}$ or $\mathcal{F}(f)$ the Fourier transform of f for both spatial and time variables,

$$\widehat{f}(\xi, \tau) = \int_{\mathbb{R}^2} e^{-ix\xi} e^{-it\tau} f(x, t) dx dt.$$

We denote by $\mathcal{F}_x$ the Fourier transform on spatial variable and if there is no confusion, we still write $\mathcal{F} = \mathcal{F}_x$. Let $\mathbb{Z}$ and $\mathbb{N}$ be the sets of integers and natural numbers, respectively. $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. For $k \in \mathbb{Z}_+$ let

$$I_k = \{\xi : |\xi| \in [2^{k-1}, 2^{k+1}]\}, \quad k \geq 1; \quad I_0 = \{\xi : |\xi| \leq 2\}.$$

Let $\eta_0 : \mathbb{R} \rightarrow [0, 1]$ denote an even smooth function supported in $[-8/5, 8/5]$ and equal to 1 in $[-5/4, 5/4]$. For $k \in \mathbb{Z}$ let $\eta_k(\xi) = \eta_0(\xi/2^k) - \eta_0(\xi/2^{k-1})$ if $k \geq 1$ and $\eta_k(\xi) \equiv 0$ if $k \leq -1$, and let $\chi_k(\xi) = \eta_0(\xi/2^k) - \eta_0(\xi/2^{k-1})$. For $k \in \mathbb{Z}$ let P_k denote the operator on $L^2(\mathbb{R})$ defined by

$$\widehat{P_k u}(\xi) = \eta_k(\xi) \widehat{u}(\xi).$$

By a slight abuse of notation we also define the operator P_k on $L^2(\mathbb{R} \times \mathbb{R})$ by the formula $\mathcal{F}(P_k u)(\xi, \tau) = \eta_k(\xi) \mathcal{F}(u)(\xi, \tau)$. For $l \in \mathbb{Z}$ let

$$P_{\leq l} = \sum_{k \leq l} P_k, \quad P_{\geq l} = \sum_{k \geq l} P_k.$$

Thus we see that $P_{\leq 0} = P_0$.

For $\phi \in \mathcal{S}'(\mathbb{R})$, we denote by $V(t)\phi = e^{-t\partial_x^3}\phi$ the free solution of linear Airy equation which is defined as

$$\mathcal{F}_x(V(t)\phi)(\xi) = \exp[i\xi^3 t] \widehat{\phi}(\xi), \quad \forall t \in \mathbb{R},$$

denote by $U_\lambda(t)\phi = e^{it\lambda\partial_x^2}\phi$ for $0 < \lambda \leq 1$ the free solution of scaled linear Schrödinger equation which is defined as

$$\mathcal{F}_x(U_\lambda(t)\phi)(\xi) = \exp[-i\lambda\xi^2 t] \widehat{\phi}(\xi), \quad \forall t \in \mathbb{R}.$$

We define the Lebesgue spaces $L_{t \in I}^q L_x^p$ and $L_x^p L_{t \in I}^q$ by the norms

$$\|f\|_{L_{t \in I}^q L_x^p} = \left\| \|f\|_{L_x^p} \right\|_{L_t^q(I)}, \quad \|f\|_{L_x^p L_{t \in I}^q} = \left\| \|f\|_{L_t^q(I)} \right\|_{L_x^p}. \quad (1.6)$$

If $I = \mathbb{R}$ we simply write $L_t^q L_x^p$ and $L_x^p L_t^q$.

We will make use of the $X^{s,b}$ -type space. Generally, let $h(\xi)$ be a continuous function, and we define

$$\|v\|_{X_{\tau=h(\xi)}^{s,b}} = \|\langle \tau - h(\xi) \rangle^b \langle \xi \rangle^s \widehat{v}(\xi, \tau)\|_{L^2(\mathbb{R}^2)}$$

where $\langle \cdot \rangle = (1 + |\cdot|^2)^{1/2}$. This type of space was first systematically studied by Bourgain [3]. In applications we usually apply $X^{s,b}$ space for b close to $1/2$. In the case $b = 1/2$ one has a good substitute: Besov-type $X^{s,b}$ space which was first noted by Tataru [20]. For $k \in \mathbb{Z}_+$ we define the frequency dyadically localized $X^{s,b}$ -type normed spaces $Y_{\tau=h(\xi)}^k$:

$$Y_{\tau=h(\xi)}^k = \left\{ f \in L^2(\mathbb{R}^2) : \begin{array}{l} f(\xi, \tau) \text{ is supported in } I_k \times \mathbb{R} \text{ and} \\ \|f\|_{Y_{\tau=h(\xi)}^k} = \sum_{j=0}^{\infty} 2^{j/2} \|\eta_j(\tau - h(\xi)) \cdot f\|_{L^2} \end{array} \right\}. \quad (1.7)$$

Then we define the l^1 -analogue of $X^{s,b}$ -type space $F_{\tau=h(\xi)}^s$ by

$$\|u\|_{F_{\tau=h(\xi)}^s}^2 = \sum_{k \geq 0} 2^{2sk} \|\eta_k(\xi) \mathcal{F}(u)\|_{Y_{\tau=h(\xi)}^k}^2. \quad (1.8)$$

In this paper we will use the space $X_{\tau=-\lambda\xi^2}^{s,b}$ and $F_{\tau=\xi^3}^s$. In order to avoid some logarithmic divergence, we use the following weaker norm for the low frequency of the KdV equation as in [10],

$$\|u\|_{\bar{Y}^0} = \|u\|_{L_x^2 L_t^\infty}.$$

For $-3/4 \leq s \leq 0$, we define

$$\bar{F}_{\tau=\xi^3}^s = \{u \in \mathcal{S}'(\mathbb{R}^2) : \|u\|_{\bar{F}_{\tau=\xi^3}^s}^2 = \sum_{k \geq 1} 2^{2sk} \|\eta_k(\xi) \mathcal{F}(u)\|_{Y_{\tau=\xi^3}^k}^2 + \|P_{\leq 0}(u)\|_{\bar{Y}^0}^2 < \infty\}.$$

For $T \geq 0$, we define the time-localized spaces $\bar{F}_{\tau=\xi^3}^s(T)$:

$$\|u\|_{\bar{F}_{\tau=\xi^3}^s(T)} = \inf_{w \in \bar{F}_{\tau=\xi^3}^s} \{ \|P_{\leq 0} u\|_{L_x^2 L_{|t| \leq T}^\infty} + \|P_{\geq 1} w\|_{\bar{F}_{\tau=\xi^3}^s} : w(t) = u(t) \text{ on } [-T, T] \}.$$

Similarly we define $X_{\tau=h(\xi)}^{s,b}(T)$.

2. L^2 BILINEAR ESTIMATES

In this section we prove some L^2 bilinear estimates which will be used to prove bilinear estimates for the coupled terms. For $\xi_1, \xi_2 \in \mathbb{R}$ let

$$\Omega_1(\xi_1, \xi_2) = -\lambda \xi_1^2 + \xi_2^3 + \lambda(\xi_1 + \xi_2)^2, \quad (2.1)$$

$$\Omega_2(\xi_1, \xi_2) = -\lambda \xi_1^2 + \lambda \xi_2^2 - (\xi_1 + \xi_2)^3. \quad (2.2)$$

Ω_1 is the resonance function for the coupled term uv , and Ω_2 is the one for $\partial_x(|u|^2)$. For compactly supported nonnegative functions $f, g, h \in L^2(\mathbb{R} \times \mathbb{R})$ we define for $m = 1, 2$

$$J_m(f, g, h) = \int_{\mathbb{R}^4} f(\xi_1, \mu_1) g(\xi_2, \mu_2) h(\xi_1 + \xi_2, \mu_1 + \mu_2 + \Omega_m(\xi_1, \xi_2)) d\xi_1 d\xi_2 d\mu_1 d\mu_2.$$

For $k, j \in \mathbb{Z}_+$ we define

$$D_{k,j}^\lambda = \{(\xi, \tau) : \xi \in I_k, \tau + \lambda \xi^2 \in I_j\},$$

and for $k \in \mathbb{Z}, j \in \mathbb{Z}_+$ we define

$$B_{k,j} = \{(\xi, \tau) : |\xi| \in [2^{k-1}, 2^{k+1}], \tau - \xi^3 \in I_j\}.$$

Let $a_1, a_2, a_3 \in \mathbb{R}$. It will be convenient to define the quantities $a_{max} \geq a_{med} \geq a_{min}$ to be the maximum, median, and minimum of a_1, a_2, a_3 respectively. Usually we use k_1, k_2, k_3 and j_1, j_2, j_3 to denote integers, $N_i = 2^{k_i}$ and $L_i = 2^{j_i}$ for $i = 1, 2, 3$ to denote dyadic numbers.

We prove the following lemma.

Lemma 2.1. *Assume $k_i \in \mathbb{Z}$, $j_i \in \mathbb{Z}_+$, and $f_{k_i, j_i} \in L^2(\mathbb{R} \times \mathbb{R})$ are nonnegative functions supported in $[2^{k_i-1}, 2^{k_i+1}] \times I_{j_i}$, $i = 1, 2, 3$. Then*

(a) *For any $k_1, k_2, k_3 \in \mathbb{Z}$ and $j_1, j_2, j_3 \in \mathbb{Z}_+$, $m = 1, 2$,*

$$J_m(f_{k_1, j_1}, f_{k_2, j_2}, f_{k_3, j_3}) \leq C 2^{j_{\min}/2} 2^{k_{\min}/2} \prod_{i=1}^3 \|f_{k_i, j_i}\|_{L^2}. \quad (2.3)$$

(b) *If $k_2 \geq 3$, then*

$$J_1(f_{k_1, j_1}, f_{k_2, j_2}, f_{k_3, j_3}) \leq C 2^{(j_{\text{med}} + j_{\text{max}})/2} 2^{-k_2} \prod_{i=1}^3 \|f_{k_i, j_i}\|_{L^2}. \quad (2.4)$$

(c) *For any $k_1, k_2, k_3 \in \mathbb{Z}$ and $j_1, j_2, j_3 \in \mathbb{Z}_+$,*

$$J_2(f_{k_1, j_1}, f_{k_2, j_2}, f_{k_3, j_3}) \leq C \lambda^{-1/2} 2^{(j_1 + j_2)/2} 2^{-k_3/2} \prod_{i=1}^3 \|f_{k_i, j_i}\|_{L^2}. \quad (2.5)$$

Proof. Let $A_{k_i}(\xi) = [\int_{\mathbb{R}} |f_{k_i, j_i}(\xi, \mu)|^2 d\mu]^{1/2}$, $i = 1, 2, 3$. Using the Cauchy-Schwartz inequality and the support properties of the functions f_{k_i, j_i} , we obtain

$$\begin{aligned} J_m(f_{k_1, j_1}, f_{k_2, j_2}, f_{k_3, j_3}) &\lesssim 2^{j_{\min}/2} \int_{\mathbb{R}^2} A_{k_1}(\xi_1) A_{k_2}(\xi_2) A_{k_3}(\xi_1 + \xi_2) d\xi_1 d\xi_2 \\ &\lesssim 2^{k_{\min}/2} 2^{j_{\min}/2} \prod_{i=1}^3 \|f_{k_i, j_i}\|_{L^2}. \end{aligned}$$

which is part (a), as desired.

For part (b), in view of the support properties of the functions, it is easy to see that $J_1(f_{k_1, j_1}, f_{k_2, j_2}, f_{k_3, j_3}) \equiv 0$ unless

$$|k_{\text{max}} - k_{\text{med}}| \leq 5, \quad 2^{j_{\text{max}}} \gtrsim |\Omega_1|. \quad (2.6)$$

We define two sets

$$A = \{(\xi_1, \xi_2) \in \mathbb{R}^2 : |2\lambda\xi_1 - 3\xi_2^2| \geq \frac{1}{2}|\xi_2|^2\}$$

and

$$B = \{(\xi_1, \xi_2) : |\Omega_1(\xi_1, \xi_2)| \geq \frac{1}{2}|\xi_2|^3\} = \{(\xi_1, \xi_2) : |\xi_2^2 + 2\lambda\xi_1 + \lambda\xi_2| \geq \frac{1}{2}|\xi_2|^2\},$$

since $|\Omega_1(\xi_1, \xi_2)| = |\xi_2(\xi_2^2 + 2\lambda\xi_1 + \lambda\xi_2)|$. We claim that

$$[2^{k_1-1}, 2^{k_1+1}] \times [2^{k_2-1}, 2^{k_2+1}] \subset A \cup B.$$

Indeed, if $(\xi_1, \xi_2) \notin A \cup B$, then

$$|4\xi_2^2 + \lambda\xi_2| \leq |3\xi_2^2 - 2\lambda\xi_1| + |\xi_2^2 + 2\lambda\xi_1 + \lambda\xi_2| \leq |\xi_2|^2,$$

which is a contradiction since $|\xi_2| \geq 2$ and $0 < \lambda \leq 1$.

For simplicity of notations we set $f_i = f_{k_i, j_i}$, $i = 1, 2, 3$. Then we get

$$\begin{aligned} & J_1(f_1, f_2, f_3) \\ & \lesssim \int_{\mathbb{R}^2} \int_A f_1(\xi_1, \mu_1) f_2(\xi_2, \mu_2) f_3(\xi_1 + \xi_2, \mu_1 + \mu_2 + \Omega_m(\xi_1, \xi_2)) d\xi_1 d\xi_2 d\mu_1 d\mu_2 \\ & \quad + \int_{\mathbb{R}^2} \int_B f_1(\xi_1, \mu_1) f_2(\xi_2, \mu_2) f_3(\xi_1 + \xi_2, \mu_1 + \mu_2 + \Omega_m(\xi_1, \xi_2)) d\xi_1 d\xi_2 d\mu_1 d\mu_2 \\ & = I + II. \end{aligned}$$

We consider first the contribution of the term I . Since for $(\xi_1, \xi_2) \in A$, then

$$|\partial_{\xi_1} \Omega_1 - \partial_{\xi_2} \Omega_1| = |2\lambda\xi_1 - 3\xi_2^2| \geq \frac{1}{2}|\xi_2|^2.$$

We will prove that if $g_i : \mathbb{R} \rightarrow \mathbb{R}_+$ are L^2 functions supported in $[2^{k_i-1}, 2^{k_i+1}]$, $i = 1, 2$, and $g : \mathbb{R}^2 \rightarrow \mathbb{R}_+$ is an L^2 function supported in $[2^{k_3-1}, 2^{k_3+1}] \times I_{j_{max}}$, then

$$\int_A g_1(\xi_1) g_2(\xi_2) g(\xi_1 + \xi_2, \Omega_1(\xi_1, \xi_2)) d\xi_1 d\xi_2 \lesssim 2^{-k_2} \|g_1\|_{L^2} \|g_2\|_{L^2} \|g\|_{L^2}. \quad (2.7)$$

This suffices for (2.4) by Cauchy-Schwartz inequality. To prove (2.7) we get

$$\begin{aligned} & \int_A g_1(\xi_1) g_2(\xi_2) g(\xi_1 + \xi_2, \Omega_1(\xi_1, \xi_2)) d\xi_1 d\xi_2 \\ & \lesssim \|g_1\|_{L^2} \|g_2\|_{L^2} \|1_A(\xi_1, \xi_2) g(\xi_1 + \xi_2, \Omega_1(\xi_1, \xi_2))\|_{L^2_{\xi_1 \xi_2}} \\ & \lesssim 2^{-k_2} \|g_1\|_{L^2} \|g_2\|_{L^2} \|g\|_{L^2}, \end{aligned}$$

where in the last inequality we used the change of the variables $x = \xi_1 + \xi_2, y = \Omega_1(\xi_1, \xi_2)$, since the Jacobi is $|\partial_{\xi_1} \Omega_1 - \partial_{\xi_2} \Omega_1| \geq \frac{1}{2}|\xi_2|^2 \gtrsim 2^{2k_2}$ in A .

Now we consider the term II . When $(\xi_1, \xi_2) \in B$, we have that $|\Omega_1| \geq \frac{1}{2}|\xi_2|^3$, so from the support properties we get $j_{max} \geq 3k_2 - 20$. Then from (a) we have in this case

$$II \lesssim 2^{j_{min}/2} 2^{k_2/2} \prod_{i=1}^3 \|f_{k_i, j_i}\|_{L^2} \lesssim 2^{j_{min}/2} 2^{j_{max}/2} 2^{-k_2} \prod_{i=1}^3 \|f_{k_i, j_i}\|_{L^2},$$

which completes the proof of (b).

Now we prove part (c). We will prove that if $g_i : \mathbb{R} \rightarrow \mathbb{R}_+$ are L^2 functions supported in $[2^{k_i-1}, 2^{k_i+1}]$, $i = 1, 2$, and $g : \mathbb{R}^2 \rightarrow \mathbb{R}_+$ is an L^2 function supported in $[2^{k_3-1}, 2^{k_3+1}] \times I_{j_{max}}$, then

$$\int_{\mathbb{R}^2} g_1(\xi_1) g_2(\xi_2) g(\xi_1 + \xi_2, \Omega_2(\xi_1, \xi_2)) d\xi_1 d\xi_2 \lesssim \lambda^{-1/2} 2^{-k_3/2} \|g_1\|_{L^2} \|g_2\|_{L^2} \|g\|_{L^2}. \quad (2.8)$$

This suffices for (2.5) by Cauchy-Schwartz inequality. To prove (2.8), we notice that

$$|\partial_{\xi_1} \Omega_2 - \partial_{\xi_2} \Omega_2| = 2\lambda|\xi_1 + \xi_2| \sim 2\lambda 2^{k_3},$$

thus by change of variable $\mu_1 = \xi_1 + \xi_2, \mu_2 = \Omega_2(\xi_1, \xi_2)$ we get

$$\|g(\xi_1 + \xi_2, \Omega_2(\xi_1, \xi_2))\|_{L^2_{\xi_1 \xi_2}} \lesssim \lambda^{-1/2} 2^{-k_3/2} \|g\|_2,$$

which is sufficient for (2.8) by Cauchy-Schwartz inequality. ■

Remark 2.2. It is easy to see from the proof that Part (a) and Part (b) of Lemma 2.1 also hold if we assume instead f_{k_i, j_i} is supported in $I_{k_i} \times I_{j_i}$ for $k_1, k_2, k_3 \in \mathbb{Z}_+$. Part (c) of Lemma 2.1 also holds if we assume instead $f_{k_1, j_1}, f_{k_2, j_2}$ are supported in $I_{k_1} \times I_{j_1}, I_{k_2} \times I_{j_2}$ respectively, for $k_1, k_2 \in \mathbb{Z}_+$.

We restate now Lemma 2.1 in a form that is suitable for the bilinear estimates in the next section.

Corollary 2.3. (a) Let $k_1, k_2, k_3, j_1, j_2, j_3 \in \mathbb{Z}_+$. Assume $f_{k_1, j_1}, f_{k_2, j_2} \in L^2(\mathbb{R} \times \mathbb{R})$ are nonnegative functions that are supported in $\{(\xi, \tau) : \xi \in I_{k_1}, \tau + \lambda \xi^2 \in I_{j_1}\}$ and $\{(\xi, \tau) : \xi \in I_{k_2}, \tau - \lambda \xi^2 \in I_{j_2}\}$ respectively, then

$$\|1_{D_{k_3, j_3}^\lambda}(\xi, \tau)(f_{k_1, j_1} * f_{k_2, j_2})\|_{L^2} \lesssim 2^{k_{\min}/2} 2^{j_{\min}/2} \prod_{i=1}^2 \|f_{k_i, j_i}\|_{L^2}. \quad (2.9)$$

Furthermore, if $k_2 > 2$, then we have

$$\|1_{D_{k_3, j_3}^\lambda}(\xi, \tau)(f_{k_1, j_1} * f_{k_2, j_2})\|_{L^2} \lesssim 2^{(j_{\min} + j_{\max})/2} 2^{-k_2} \prod_{i=1}^2 \|f_{k_i, j_i}\|_{L^2}. \quad (2.10)$$

(b) Let $k_1, k_2, j_1, j_2, j_3 \in \mathbb{Z}_+$ and $k_3 \in \mathbb{Z}$. Assume $f_{k_1, j_1}, f_{k_2, j_2} \in L^2(\mathbb{R} \times \mathbb{R})$ are nonnegative functions that are supported in $\{(\xi, \tau) : \xi \in I_{k_1}, \tau + \lambda \xi^2 \in I_{j_1}\}$ and $\{(\xi, \tau) : \xi \in I_{k_2}, \tau - \lambda \xi^2 \in I_{j_2}\}$ respectively, then

$$\|1_{B_{k_3, j_3}}(\xi, \tau)(f_{k_1, j_1} * f_{k_2, j_2})\|_{L^2} \lesssim 2^{k_{\min}/2} 2^{j_{\min}/2} \prod_{i=1}^2 \|f_{k_i, j_i}\|_{L^2}, \quad (2.11)$$

and also

$$\|1_{B_{k_3, j_3}}(\xi, \tau)(f_{k_1, j_1} * f_{k_2, j_2})\|_{L^2} \lesssim \lambda^{-1/2} 2^{j_1/2} 2^{j_2/2} 2^{-k_3/2} \prod_{i=1}^2 \|f_{k_i, j_i}\|_{L^2}. \quad (2.12)$$

Proof. We just prove (a), the proof for (b) is similar. Clearly, we have

$$\|1_{D_{k_3, j_3}^\lambda}(\xi, \tau)(f_{k_1, j_1} * f_{k_2, j_2})(\xi, \tau)\|_{L^2} = \sup_{\|f\|_{L^2}=1} \left| \int_{D_{k_3, j_3}^\lambda} f \cdot f_{k_1, j_1} * f_{k_2, j_2} d\xi d\tau \right|.$$

We denote $f_{k_3, j_3} = 1_{D_{k_3, j_3}^\lambda}(\xi, \tau) \cdot f(\xi, \tau)$. Define $f_{k_1, j_1}^\#(\xi, \mu) = f_{k_1, j_1}(\xi, \mu - \lambda \xi^2)$, $f_{k_2, j_2}^\#(\xi, \mu) = f_{k_2, j_2}(\xi, \mu + \lambda \xi^2)$, $f_{k_3, j_3}^\#(\xi, \mu) = f_{k_3, j_3}(\xi, \mu - \lambda \xi^2)$. Then for $i = 1, 2, 3$ the functions $f_{k_i, j_i}^\#$ are supported in $I_{k_i} \times I_{j_i}$ and $\|f_{k_i, j_i}^\#\|_{L^2} = \|f_{k_i, j_i}\|_{L^2}$. Using simple changes of variables, we get that

$$\int_{D_{k_3, j_3}^\lambda} f \cdot f_{k_1, j_1} * f_{k_2, j_2} d\xi d\tau = J_1(f_{k_1, j_1}^\#, f_{k_2, j_2}^\#, f_{k_3, j_3}^\#).$$

Then (a) follows from Lemma 2.1 (a), (b) and Remark 2.2. ■

3. BILINEAR ESTIMATES FOR THE COUPLING TERMS

This section is devoted to prove the bilinear estimates for the coupling terms in the F^s -type space. We first recall an abstract extension lemma. For a continuous function $h(\xi)$, define group $W_h(t)$ by

$$W_h(t)f = \mathcal{F}_x e^{ith(\xi)} \mathcal{F}_x f.$$

Then $W_h(t)f$ is the solution to the following equation

$$\partial_t u - ih(-i\partial_x)u = 0, \quad u(x, 0) = f(x).$$

Lemma 3.1 (Extension lemma). *Assume h is a continuous function. Let Z be any space-time Banach space which obeys the time modulation estimate*

$$\|g(t)F(t, x)\|_Z \leq \|g\|_{L_t^\infty} \|F(t, x)\|_Z \quad (3.1)$$

for any $F \in Z$ and $g \in L_t^\infty$. Moreover, if for all $u_0 \in L_x^2$

$$\|W_h(t)u_0\|_Z \lesssim \|u_0\|_{L_x^2}.$$

Then one also has the estimate that for all $k \in \mathbb{Z}_+$ and $u \in F^0$

$$\|P_k(u)\|_Z \lesssim \|\widehat{P_k(u)}\|_{Y_{\tau=h(\xi)}^k}.$$

Proof. We refer the readers to Lemma 3.2 in [10]. ■

Proposition 3.2 ($X_{\tau=-\lambda\xi^2}^k$ embedding). *Let $k \in \mathbb{Z}_+$, $j \in \mathbb{N}$. Assume $u \in \mathcal{S}(\mathbb{R}^2)$, then we have*

$$\begin{aligned} \|P_k(u)\|_{L_t^q L_x^r} &\lesssim \lambda^{-1/q} \|P_k(u)\|_{Y_{\tau=-\lambda\xi^2}^k}, \\ \|P_j(u)\|_{L_x^\infty L_t^2} &\lesssim \lambda^{-1/2} 2^{-j/2} \|\mathcal{F}[P_j(u)]\|_{Y_{\tau=-\lambda\xi^2}^k}, \end{aligned}$$

where (q, r) satisfies $2 \leq q, r \leq \infty$ and $2/q = 1/2 - 1/r$. As a consequence, we get from the definition that for $u \in F_{\tau=-\lambda\xi^2}^s$

$$\|u\|_{L_t^\infty H^s} \lesssim \|u\|_{F_{\tau=-\lambda\xi^2}^s}.$$

Proof. From Lemma 3.1, it suffices to prove

$$\begin{aligned} \|U_\lambda(t)f\|_{L_t^q L_x^r} &\lesssim \lambda^{-1/q} \|f\|_{L_x^2}, \\ \|U_\lambda(t)f\|_{L_x^\infty L_t^2} &\lesssim \lambda^{-1/2} \|f\|_{\dot{H}_x^{-1/2}}, \end{aligned}$$

which are well-known, for example see [13] and [14]. ■

Proposition 3.3 ($X_{\tau=\xi^3}^k$ embedding). *Let $k \in \mathbb{Z}_+$, $j \in \mathbb{N}$. Assume $u \in F^0$, then we have*

$$\|P_k(u)\|_{L_t^q L_x^r} \lesssim \|P_k(u)\|_{Y_{\tau=\xi^3}^k}, \quad (3.2)$$

$$\|P_k(u)\|_{L_x^2 L_{t \in I}^\infty} \lesssim 2^{3k/4} \|\mathcal{F}[P_k(u)]\|_{Y_{\tau=\xi^3}^k}, \quad (3.3)$$

$$\|P_j(u)\|_{L_x^\infty L_t^2} \lesssim 2^{-j} \|\mathcal{F}[P_j(u)]\|_{Y_{\tau=\xi^3}^j}, \quad (3.4)$$

where (q, r) satisfies $2 \leq q, r \leq \infty$ and $3/q = 1/2 - 1/r$. As a consequence, we get from the definition that for $u \in \bar{F}_{\tau=\xi^3}^s$, then

$$\|u\|_{L_t^\infty H^s} \lesssim \|u\|_{\bar{F}_{\tau=\xi^3}^s}.$$

Proof. We refer the readers to proposition 3.3 in [10]. ■

The main result of this section is the following lemma.

Lemma 3.4. (a) If $s_1 \geq 0$, $s_2 \in (-1, -1/2)$, $s_1 - s_2 < 1$, $0 < \theta \ll 1$, and $u \in X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}$, $v \in \bar{F}_{\tau=\xi^3}^{s_2}$ then

$$\|\psi(t)uv\|_{X_{\tau=-\lambda\xi^2}^{s_1, -1/2+2\theta}} \lesssim \lambda^{-1/2} \|u\|_{X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}} \|v\|_{\bar{F}_{\tau=\xi^3}^{s_2}}. \quad (3.5)$$

(b) If $s_1 \geq 0$, $s_2 - s_1 < -1/2$, and $u, w \in X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}$ then

$$\|\psi(t)\partial_x(u\bar{w})\|_{X_{\tau=\xi^3}^{s_2, -1/2+2\theta}} \lesssim \lambda^{-1/2} \|u\|_{X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}} \|w\|_{X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}}. \quad (3.6)$$

Proof. We first prove (a). From the definition, we get that the left-hand side of (3.5) is dominated by

$$\left\| 2^{s_1 k_3} \sum_{j_3=0}^{\infty} 2^{-1/2 j_3 + 2\theta j_3} \|1_{D_{k_3, j_3}^\lambda} \cdot \widehat{\psi(t)uv}\|_{L_{\xi, \tau}^2} \right\|_{l_{k_3}^2}. \quad (3.7)$$

Now we begin to estimate

$$\sum_{j_3}^{\infty} 2^{-1/2 j_3 + 2\theta j_3} \|1_{D_{k_3, j_3}^\lambda} \cdot \widehat{\psi(t)uv}\|_{L_{\xi, \tau}^2}. \quad (3.8)$$

Decomposing u, v , for $k_1, j_1 \in \mathbb{Z}_+$ set

$$\begin{aligned} f_{k_1, j_1}(\xi_1, \tau_1) &= \eta_{k_1}(\xi_1) \eta_{j_1}(\tau_1 + \lambda \xi_1^2) \widehat{\psi(t/2)u}(\xi_1, \tau_1), \\ f_{k_2, j_2}(\xi_2, \tau_2) &= \eta_{k_2}(\xi_2) \eta_{j_2}(\tau_2 - \xi_2^3) \widehat{\psi(t/2)v}(\xi_2, \tau_2), \end{aligned}$$

then we get for fixed k_3 ,

$$(3.8) \lesssim \sum_{(k_1, k_2, k_3) \in K} \sum_{j_3 \geq 0} 2^{-j_3/2 + 2\theta j_3} \sum_{j_1, j_2 \geq 0} \|1_{D_{k_3, j_3}^\lambda} \cdot f_{k_1, j_1} * f_{k_2, j_2}\|_{L_{\xi_3 \tau_3}^2}. \quad (3.9)$$

where $K = \{(k_1, k_2, k_3) \in \mathbb{Z}_+^3 : |k_{med} - k_{max}| < 5\}$, since it is easy to see that $1_{D_{k_3, j_3}^\lambda} \cdot f_{k_1, j_1} * f_{k_2, j_2} \equiv 0$ unless

$$|k_{med} - k_{max}| < 5.$$

We may also assume that $k_{max} \geq 20$, since for $k_{max} \leq 20$ we can get from Plancherel's equality that

$$\begin{aligned} (3.8) &\lesssim \sum_{j_3 \geq 0} 2^{-j_3/2 + 2\theta j_3} \|1_{D_{k_3, j_3}^\lambda} \cdot \widehat{P_{\leq 20} u} * \widehat{\psi(t)P_{\leq 20} v}\|_{L_{\xi_3 \tau_3}^2} \\ &\lesssim \|P_{\leq 20} u \psi(t) P_{\leq 20} v\|_{L_{x, t}^2} \\ &\lesssim \|P_{\leq 20} u\|_{L_t^\infty L_x^2} \|\psi(t) P_{\leq 20} v\|_{L_x^2 L_t^\infty} \end{aligned}$$

which suffices to give the bound for this case by Proposition 3.2, 3.3.

Now we assume $k_{max} \geq 20$ and prove (3.9). First we assume that $k_2 \geq 2$. Clearly we may also assume that $j_{max} \leq 10k_2$, otherwise, we can apply (2.9), then we have a 2^{-5k_2} to spare. After these assumptions, we can make use of (2.10) to bound (3.9) by

$$\sum_{(k_1, k_2, k_3) \in K} \sum_{k_2 \geq 2} \sum_{j_3 \geq 0} 2^{-j_3/2 + 2\theta j_3} \sum_{j_1, j_2 \geq 0} 2^{j_{max}/2} 2^{j_{med}/2} 2^{-k_2} \prod_{i=1}^2 \|f_{k_i, j_i}\|_{L^2} \quad (3.10)$$

when $j_3 = j_{min}$, then (3.10) is bounded by

$$\begin{aligned} & \sum_{(k_1, k_2, k_3) \in K, k_2 \geq 2} \sum_{j_1, j_2 \geq 0} 2^{j_1/2} 2^{j_2/2} 2^{-k_2} \prod_{i=1}^2 \|f_{k_i, j_i}\|_{L^2} \\ & \lesssim \sum_{(k_1, k_2, k_3) \in K, k_2 \geq 2} 2^{-k_2} \|P_{k_1} u\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|P_{k_2} v\|_{Y_{\tau=\xi^3}^{k_2}}, \end{aligned} \quad (3.11)$$

when $j_1 = j_{min}$, then (3.10) is bounded by

$$\begin{aligned} & \sum_{(k_1, k_2, k_3) \in K, k_2 \geq 2} \sum_{j_3 \geq 0} 2^{2\theta j_3} \sum_{j_1, j_2 \geq 0} 2^{j_2/2} 2^{-k_2} \prod_{i=1}^2 \|f_{k_i, j_i}\|_{L^2} \\ & \lesssim \sum_{(k_1, k_2, k_3) \in K, k_2 \geq 2} 2^{2\theta k_2} \sum_{j_1, j_2 \geq 0} 2^{j_1/2} 2^{j_2/2} 2^{-k_2} \prod_{i=1}^2 \|f_{k_i, j_i}\|_{L^2} \\ & \lesssim \sum_{(k_1, k_2, k_3) \in K, k_2 \geq 2} 2^{-k_2+20\theta k_2} \|P_{k_1} u\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|P_{k_2} v\|_{Y_{\tau=\xi^3}^{k_2}}, \end{aligned} \quad (3.12)$$

where we use $j_{max} \leq 10k_2$ in the last step. The case $j_2 = j_{min}$ is the same as $j_1 = j_{min}$, so we omit the details.

Dividing the summation on k_1, k_2 in the right-hand side of (3.10) into several parts, we get from (3.11), (3.12)

$$(3.10) \lesssim \sum_{i=1}^5 \sum_{A_i(k_3)} 2^{-k_2+20\theta k_2} \|P_{k_1} u\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|P_{k_2} v\|_{Y_{\tau=\xi^3}^{k_2}}$$

where we denote

$$\begin{aligned} A_1(k_3) &= \{|k_2 - k_3| \leq 5, k_1 \leq k_2 - 10, \text{ and } k_2 \geq 30\}; \\ A_2(k_3) &= \{|k_1 - k_3| \leq 5, 2 \leq k_2 \leq k_1 - 10, \text{ and } k_1 \geq 30\}; \\ A_3(k_3) &= \{|k_1 - k_2| \leq 5, k_3 \leq k_2 - 10, \text{ and } k_1 \geq 30\}; \\ A_4(k_3) &= \{|k_2 - k_3| \leq 10, |k_1 - k_2| \leq 10, \text{ and } k_2 \geq 30\}; \\ A_5(k_3) &= \{k_1, k_2, k_3 \leq 200\}. \end{aligned}$$

So we can bounded (3.8) by

$$\sum_{i=1}^5 \left\| 2^{s_1 k_3} \sum_{A_i(k_3)} 2^{-k_2+20\theta k_2} \|P_{k_1} u\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|P_{k_2} v\|_{Y_{\tau=\xi^3}^{k_2}} \right\|_{l_{k_3}^2}. \quad (3.13)$$

Now we begin to estimate (3.13) case by case. For case A_1 , by Cauchy-Schwartz's inequality we have

$$\begin{aligned} & \left\| 2^{s_1 k_3} \sum_{A_1(k_3)} 2^{-k_2+20\theta k_2} \|P_{k_1} u\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|P_{k_2} v\|_{Y_{\tau=\xi^3}^{k_2}} \right\|_{l_{k_3}^2} \\ &= \left\| 2^{s_1 k_3} \sum_{|k_2 - k_3| \leq 5, k_2 \geq 20} \sum_{k_1 \leq k_2 - 10} 2^{-k_2+20\theta k_2} \|P_{k_1} u\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|P_{k_2} v\|_{Y_{\tau=\xi^3}^{k_2}} \right\|_{l_{k_3}^2} \\ &\lesssim \left\| 2^{s_1 k_3} \sum_{|k_2 - k_3| \leq 5, k_2 \geq 20} 2^{-k_2+21\theta k_2} \|P_{k_2} v\|_{Y_{\tau=\xi^3}^{k_2}} \right\|_{l_{k_3}^2} \left(\sum_{k_1 \geq 0} 2^{2s_1 k_1} \|P_{k_1} u\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \right)^{1/2}. \end{aligned} \quad (3.14)$$

since $s_1 - s_2 < 1$, $s_1 \geq 0$, $s_2 > -1$ and $0 < \theta \ll 1$, we can bound (3.14) by $\|v\|_{\bar{F}_{\tau=\xi^3}^{s_2}} \|u\|_{X_{\tau=\lambda\xi^2}^{s_1, 1/2+\theta}}$. The proof for $A_2 - A_5$ are similar, we omit here.

Now we assume that $k_2 \leq 2$, by the assumption $k_{max} \geq 20$, we have $|k_1 - k_3| \leq 5$, and $k_1, k_2 \geq 10$. For simplicity of notations we assume $k_1 = k_3 = k$, then the left hand side of (3.8) can be estimated as

$$\sum_{k \geq 0} 2^{s_1 k} \|\widehat{P_k(u)} * \widehat{\psi(t)P_{\leq 2}(v)}\|_{L^2} \lesssim \sum_{k \geq 0} 2^{s_1 k} \|P_k(u)\|_{L_x^\infty L_t^2} \|\psi(t)P_{\leq 2}(v)\|_{L_x^2 L_t^\infty}. \quad (3.15)$$

This is enough in view of the definition, Proposition 3.2 and Proposition 3.3.

Now we begin to prove (b), from the definition, we only need to prove

$$\|\partial_x(u\bar{w})\|_{X_{\tau=\xi^3}^{s_2, -1/2+2\theta}} \lesssim \lambda^{-1/2} \|u\|_{F_{\tau=-\lambda\xi^2}^{s_1}} \|w\|_{F_{\tau=-\lambda\xi^2}^{s_1}}. \quad (3.16)$$

By the definition and $\theta \ll 1$, we have

$$\begin{aligned} \|\partial_x(u\bar{w})\|_{X_{\tau=\xi^3}^{s_2, -1/2+2\theta}} &\lesssim \left(\sum_{k \geq 1} 2^{2s_2 k} \left(\sum_{j \geq 0} 2^{-j/4} \|1_{B_{k,j}}(\xi, \tau) i\xi \mathcal{F}(u\bar{w})\|_{L_{\xi, \tau}^2} \right)^2 \right)^{1/2} \\ &\quad + \left(\sum_{k \leq 0} \left(\sum_{j \geq 0} 2^{-j/4} \|1_{B_{k,j}}(\xi, \tau) i\xi \mathcal{F}(u\bar{w})\|_{L_{\xi, \tau}^2} \right)^2 \right)^{1/2} \\ &:= I + II. \end{aligned} \quad (3.17)$$

Now let

$$\begin{aligned} f_{k_1, j_1}(\xi_1, \tau_1) &= \eta_{k_1}(\xi_1) \eta_{j_1}(\tau_1 + \lambda \xi_1^2) \hat{u}(\xi_1, \tau_1), \\ \overline{g_{k_2, j_2}(\xi_2, \tau_2)} &= \eta_{k_1}(\xi_2) \eta_{j_1}(\tau_2 - \lambda \xi_2^2) \tilde{w}(\xi_2, \tau_2), \end{aligned}$$

then we have

$$\begin{aligned} \|u\|_{F_{\tau=-\lambda\xi^2}^s} &= \left\| 2^{s k_1} \sum_{j_1 \geq 0} 2^{j_1/2} \|f_{k_1, j_1}\|_{L^2} \right\|_{l_{k_1}^2}, \\ \|w\|_{F_{\tau=-\lambda\xi^2}^s} &= \left\| 2^{s k_2} \sum_{j_2 \geq 0} 2^{j_2/2} \|g_{k_2, j_2}\|_{L^2} \right\|_{l_{k_2}^2}. \end{aligned}$$

Then from the definition, I and II in (3.17) can be bounded as following

$$I \lesssim \left\| 2^{(s_2+1)k_3} \sum_{(k_1, k_2, k_3) \in K} \sum_{j_3 \geq 0} 2^{-j_3/4} \sum_{j_1, j_2 \geq 0} \|1_{B_{k_3, j_3}} f_{k_1, j_1} * g_{k_2, j_2}\|_{L_{\xi_3 \tau_3}^2} \right\|_{l_{k_3 \geq 1}^2} \quad (3.18)$$

and

$$II \lesssim \left\| 2^{k_3} \sum_{(k_1, k_2, k_3) \in K} \sum_{j_3 \geq 0} 2^{-j_3/4} \sum_{j_1, j_2 \geq 0} \|1_{B_{k_3, j_3}} f_{k_1, j_1} * g_{k_2, j_2}\|_{L_{\xi_3 \tau_3}^2} \right\|_{l_{k_3 \leq 0}^2}. \quad (3.19)$$

We first estimate term I in (3.18). By (2.12) we have

$$\begin{aligned} &\sum_{(k_1, k_2, k_3) \in K} \sum_{j_3 \geq 0} 2^{-j_3/4} \sum_{j_1, j_2 \geq 0} \|1_{B_{k_3, j_3}} f_{k_1, j_1} * g_{k_2, j_2}\|_{L_{\xi_3 \tau_3}^2} \\ &\lesssim \sum_{(k_1, k_2, k_3) \in K, k_3 \geq 1} 2^{-k_3/2} \sum_{j_1, j_2 \geq 0} \lambda^{-1/2} 2^{j_1/2} 2^{j_2/2} \|f_{k_1, j_1}\|_{L^2} \|g_{k_2, j_2}\|_{L^2} \\ &\lesssim \lambda^{-1/2} \sum_{(k_1, k_2, k_3) \in K, k_3 \geq 1} 2^{-k_3/2} \|\widehat{P_{k_1} u}\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|\widehat{P_{k_2} w}\|_{Y_{\tau=-\lambda\xi^2}^{k_2}}. \end{aligned} \quad (3.20)$$

Dividing the summation on k_1, k_2 in the right-hand side of (3.20) into several parts,

$$(3.20) \lesssim \lambda^{-1/2} \sum_{i=1}^5 \sum_{B_i(k_3)} \|P_{k_1} u\|_{F_{\tau=\lambda\xi^2}^{k_1}} \|P_{k_2} w\|_{F_{\tau=\lambda\xi^2}^{k_2}}$$

where we denote

$$\begin{aligned} B_1(k_3) &= \{|k_2 - k_3| \leq 5, k_1 \leq k_2 - 10, \text{ and } k_2 \geq 30\}; \\ B_2(k_3) &= \{|k_1 - k_3| \leq 5, k_2 \leq k_1 - 10, \text{ and } k_1 \geq 30\}; \\ B_3(k_3) &= \{|k_1 - k_2| \leq 5, 1 \leq k_3 \leq k_2 - 10, \text{ and } k_1 \geq 30\}; \\ B_4(k_3) &= \{|k_2 - k_3| \leq 10, |k_1 - k_2| \leq 10, \text{ and } k_2 \geq 30\}; \\ B_5(k_3) &= \{k_1, k_2, k_3 \leq 200, k_3 \geq 1\}. \end{aligned}$$

So we can bounded (3.18) by

$$I \lesssim \lambda^{-1/2} \sum_{i=1}^5 \left\| 2^{(s_2+1/2)k_3} \sum_{B_i(k_3)} \|\widehat{P_{k_1} u}\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|\widehat{P_{k_2} w}\|_{Y_{\tau=-\lambda\xi^2}^{k_2}} \right\|_{l_{k_3}^2}. \quad (3.21)$$

Now we begin to estimate (3.21) case by case. For case B_1 , by Cauchy-Schwartz's inequality we have

$$\begin{aligned} & \lambda^{-1/2} \left\| 2^{(s_2+1/2)k_3} \sum_{B_1(k_3)} \|\widehat{P_{k_1} u}\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \|\widehat{P_{k_2} w}\|_{Y_{\tau=-\lambda\xi^2}^{k_2}} \right\|_{l_{k_3}^2} \\ &= \lambda^{-1/2} \left\| 2^{(s_2+1/2)k_3} \sum_{|k_2-k_3| \leq 5, k_2 \geq 20} \|\widehat{P_{k_2} w}\|_{Y_{\tau=-\lambda\xi^2}^{k_2}} \sum_{k_1 \leq k_2-10} \|\widehat{P_{k_1} u}\|_{Y_{\tau=-\lambda\xi^2}^{k_1}} \right\|_{l_{k_3}^2} \\ &\lesssim \lambda^{-1/2} \left\| 2^{(s_2+1/2+\theta)k_3} \sum_{|k_2-k_3| \leq 5, k_2 \geq 20} \|\widehat{P_{k_2} w}\|_{Y_{\tau=-\lambda\xi^2}^{k_2}} \right\|_{l_{k_3}^2} \|u\|_{F_{\tau=-\lambda\xi^2}^{s_1}}. \quad (3.22) \end{aligned}$$

since $s_2 - s_1 < -1/2$, $s_1 \geq 0$ and $0 < \theta \ll 1$, so we can bounded (3.22) by $\|u\|_{X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}} \|w\|_{X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}}$. The proof for $B_2 - B_5$ are similar, we omit here.

Use the same argument as above, under the restriction $s_2 - s_1 < -1/2$, $s_1 \geq 0$ and $0 < \theta \ll 1$, we can bound the part II in (3.19) by $\|u\|_{X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}} \|w\|_{X_{\tau=-\lambda\xi^2}^{s_1, 1/2+\theta}}$.

Thus we finish the proof. $\blacksquare$

4. PROOF OF THE MAIN THEOREM

In this section we prove Theorem 1.1. We first recall some linear estimates in $X_{\tau=h(\xi)}^{s,b}$ and $\bar{F}_{\tau=h(\xi)}^s$. Let $W_h(t)f = \mathcal{F}_x^{-1} e^{ith(\xi)} \mathcal{F}_x f$. The following lemma has been proved by Kenig, Ponce and Vega in [15], and then improved by Ginibre, Y. Tsutsumi and Velo in [7].

Lemma 4.1. *Let $s \in \mathbb{R}$, $-1/2 < b' \leq 0 \leq b \leq b' + 1$ and $T \in [0, 1]$. Then for $u_0 \in H^s$ and $F \in X_{\tau=h(\xi)}^{s,b}$ we have*

$$\begin{aligned} & \|\psi_1(t) W_h(t) u_0\|_{X_{\tau=h(\xi)}^{s,b}} \lesssim \|u_0\|_{H^s}, \\ & \left\| \psi_T(t) \int_0^t W_h(t-t') F(t', \cdot) dt' \right\|_{X_{\tau=h(\xi)}^{s,b}} \lesssim T^{1-b+b'} \|F\|_{X_{\tau=h(\xi)}^{s,b'}}. \end{aligned}$$

Next we prove the linear estimates in $\bar{F}^s$. The proof was essentially given in [12] for the Benjamin-Ono equation.

Lemma 4.2. (a) Assume $s \in \mathbb{R}$ and $\phi \in H^s$. Then there exists $C > 0$ such that

$$\|\psi(t)W_h(t)\phi\|_{F_{\tau=h(\xi)}^s} \leq C\|\phi\|_{H^s}. \quad (4.1)$$

(b) Assume $s \in \mathbb{R}, k \in \mathbb{Z}_+$ and u satisfies $(i + \tau - \xi^3)^{-1}\mathcal{F}(u) \in Y_{\tau=h(\xi)}^k$. Then there exists $C > 0$ such that

$$\left\| \mathcal{F} \left[\psi(t) \int_0^t W_h(t-s)(u(s))ds \right] \right\|_{Y_{\tau=h(\xi)}^k} \leq C \|(i + \tau - \xi^3)^{-1}\mathcal{F}(u)\|_{Y_{\tau=h(\xi)}^k}. \quad (4.2)$$

Now we give the estimates for the cubic nonlinear term.

Lemma 4.3. Let $u, u' \in X_{\tau=-\lambda\xi^2}^{s,b}$ with $1/2 < b < 1$ and $s \geq 0$. Then for all $a \geq 0$ we have that

$$\begin{aligned} \| |u|^2 u \|_{X_{\tau=-\lambda\xi^2}^{s,-a}} &\lesssim \lambda^{-1/2} \|u\|_{X_{\tau=-\lambda\xi^2}^{s,b}}^3, \\ \| |u|^2 u - |u'|^2 u' \|_{X_{\tau=-\lambda\xi^2}^{s,-a}} &\lesssim \lambda^{-1/2} (\|u\|_{X_{\tau=-\lambda\xi^2}^{s,b}}^2 + \|u'\|_{X_{\tau=-\lambda\xi^2}^{s,b}}^2) \|u - u'\|_{X_{\tau=-\lambda\xi^2}^{s,b}}. \end{aligned}$$

Proof. We just proof the first one when $s = 0$ for example. By Hölder inequality and (3), we have

$$\| |u|^2 u \|_{X_{\tau=-\lambda\xi^2}^{0,-a}} \leq \| |u|^2 u \|_{L^2} = \|u\|_{L_{t,x}^6}^3 \leq C\lambda^{-1/2} \|u\|_{X_{\tau=-\lambda\xi^2}^{0,b}}^3$$

Thus we finish the proof. $\blacksquare$

We also need a bilinear estimates. For $u, v \in \bar{F}^s$ we define the bilinear operator

$$B(u, v) = \psi(t/4) \int_0^t W(t-\tau) \partial_x (\psi^2(\tau) u(\tau) \cdot v(\tau)) d\tau.$$

The following lemma is due to the first author [10], which is the key to get the global well-posedness for KdV in $H^{-3/4}$.

Lemma 4.4 (Proposition 4.2, [10]). Assume $-3/4 \leq s \leq 0$. Then there exists $C > 0$ such that

$$\|B(u, v)\|_{\bar{F}^s} \leq C(\|u\|_{\bar{F}^s} \|v\|_{\bar{F}^{-3/4}} + \|u\|_{\bar{F}^{-3/4}} \|v\|_{\bar{F}^s}) \quad (4.3)$$

hold for any $u, v \in \bar{F}^s$.

Now we are ready to prove Theorem 1.1. We consider first (1.4) under the condition (1.5). We may assume $\alpha = \beta = \gamma = 1$. From Duhamel's principle, the Cauchy problem (1.4) is equivalent to the following integral equation system

$$\begin{aligned} u(t) &= U_\lambda(t)\phi_1(x) - i \int_0^t U_\lambda(t-t') [\lambda(uv)(t') + \lambda^{-1}(|u|^2 u)(t')] dt' \\ v(t) &= V(t)\phi_2(x) + \int_0^t V(t-t') \left[-\frac{1}{2} \partial_x(v^2)(t') + \partial_x(|u|^2)(t') \right] dt'. \end{aligned} \quad (4.4)$$

To study the local existence, it suffices to study the following time localized system

$$\begin{aligned} u(t) &= \psi_1(t)U_\lambda(t)\phi_1(x) - i\psi_T(t) \int_0^t U_\lambda(t-t') [\lambda(uv)(t') + \lambda^{-1}(|u|^2 u)(t')] dt' \\ v(t) &= \psi_1(t)V(t)\phi_2(x) - \frac{1}{2}\psi_4(t) \int_0^t V(t-t') \partial_x(v^2)(t') dt' \\ &\quad + \psi_T(t) \int_0^t V(t-t') \partial_x(|u|^2)(t') dt'. \end{aligned} \quad (4.5)$$

It is easy to see that if (u, v) solve the system (4.5) for all $t \in \mathbb{R}$, then (u, v) also solve the system (4.4) for $|t| \leq T$.

We follow the similar argument as the one given in [6] to construct our solution spaces. We consider the following function space where we seek our solution:

$$\Sigma_\theta := \left\{ (u, v) \in X_{\tau=-\lambda\xi^2}^{0,1/2+\theta} \times \bar{F}_{\tau=\xi^3}^{-3/4}; \quad \|u\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}} \leq M \text{ and } \|v\|_{\bar{F}_{\tau=\xi^3}^{-3/4}} \leq \epsilon_0 \right\},$$

where $0 < \theta, \epsilon_0 \ll 1$ and $M > 0$, will be chosen later. Furthermore, Σ_θ is a complete metric space with norm

$$\|(u, v)\|_{\Sigma_\theta} = \|u\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}} + \|v\|_{\bar{F}_{\tau=\xi^3}^{-3/4}}.$$

For $(u, v) \in \Sigma_\theta$, we define the maps $\Phi = \Phi_1 \times \Phi_2$

$$\Phi_1(u, v) = \psi_1(t)U_\lambda(t)\phi_1(x) - i\psi_T(t) \int_0^t U_\lambda(t-t')[\lambda(uv)(t') + \lambda^{-1}(|u|^2u)(t')]dt'$$

and

$$\begin{aligned} \Phi_2(u, v) = & \psi_1(t)V(t)\phi_2(x) - \frac{1}{2}\psi_4(t) \int_0^t V(t-t')\partial_x(v^2)(t')dt' \\ & + \psi_T(t) \int_0^t V(t-t')\partial_x(|u|^2)(t')dt'. \end{aligned}$$

Then from Lemma 4.1-4.4, we get

$$\begin{aligned} \|\Phi_1(u, v)\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}} & \leq c_0\|\phi_1\|_{L^2} + c_1T^\theta \left[\lambda\|uv\|_{X_{\tau=-\lambda\xi^2}^{0,-1/2+2\theta}} + \lambda^{-1}\||u|^2u\|_{X_{\tau=-\lambda\xi^2}^{0,-1/2+2\theta}} \right] \\ & \leq c_0\|\phi_1\|_{L^2} + c_1T^\theta \left[\lambda\|u\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}}\|v\|_{\bar{F}_{\tau=\xi^3}^{-3/4}} + \lambda^{-3/2}\|u\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}}^3 \right] \\ & \leq c_0\|\phi_1\|_{L^2} + c_1T^\theta [\lambda M\epsilon_0 + \lambda^{-2}M^3], \end{aligned}$$

and

$$\begin{aligned} \|\Phi_2(u, v)\|_{\bar{F}_{\tau=\xi^3}^{-3/4}} & \leq c_0\|\phi_2\|_{H^{-3/4}} + \frac{1}{2}\left\|\psi\left(\frac{t}{4}\right) \int_0^t V(t-t')\partial_x(v^2)(t')dt'\right\|_{\bar{F}_{\tau=\xi^3}^{-3/4}} \\ & \quad + \left\|\psi_T(t) \int_0^t V(t-t')\partial_x(|u|^2)(t')dt'\right\|_{X_{\tau=\xi^3}^{-3/4,1/2+\theta}} \\ & \leq c_0\|\phi_2\|_{H^{-3/4}} + c_2\|v\|_{\bar{F}_{\tau=\xi^3}^{-3/4}}^2 + c_2T^\theta\|\partial_x|u|^2\|_{X_{\tau=\xi^3}^{-3/4,-1/2+2\theta}} \\ & \leq c_0\|\phi_2\|_{H^{-3/4}} + c_2\|v\|_{\bar{F}_{\tau=\xi^3}^{-3/4}}^2 + c_2\lambda^{-1/2}T^\theta\|u\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}}^2 \\ & \leq c_0\|\phi_2\|_{H^{-3/4}} + c_2\epsilon_0^2 + c_2\lambda^{-1/2}T^\theta M^2. \end{aligned}$$

Now taking $M = 2c_0\|\phi_1\|_{L^2}$ and $\epsilon_0 \ll 1$ satisfy $c_2\epsilon_0 = 1/8$, then we have that

$$\begin{aligned} \|\Phi_1(u, v)\|_{X_\lambda^{0,1/2+\theta}} & \leq \frac{M}{2} + c_1T^\theta [\lambda M\epsilon_0 + \lambda^{-3/2}M^3], \\ \|\Phi_2(u, v)\|_{\bar{F}^{-3/4}} & \leq \frac{\epsilon_0}{2} + c_2\epsilon_0^2 + c_2\lambda^{-1/2}T^\theta M^2. \end{aligned}$$

Therefore, we choose $T > 0$ such that

$$T^\theta \ll \frac{\lambda^{1/2}}{M^2} \min(\epsilon_0, \lambda), \quad (4.6)$$

then $(\Phi_1(u, v), \Phi_2(u, v)) \in \Sigma_\theta$. Similarly we have that

$$\begin{aligned} \|\Phi_1(u, v) - \Phi_1(u', v')\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}} &\leq c_3 \lambda^{-3/2} T^\theta (M + \epsilon_0 + M^2) \\ &\quad \cdot [\|u - u'\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}} + \|v - v'\|_{\bar{F}_{\tau=\xi^3}^{-3/4}}], \\ \|\Phi_2(u, v) - \Phi_2(u', v')\|_{\bar{F}_{\tau=\xi^3}^{-3/4}} &\leq c_3 (\lambda^{-1/2} T^\theta M + \epsilon_0) \\ &\quad \cdot [\|u - u'\|_{X_{\tau=-\lambda\xi^2}^{0,1/2+\theta}} + \|v - v'\|_{\bar{F}_{\tau=\xi^3}^{-3/4}}]. \end{aligned}$$

Thus for T in (4.6) we get

$$\|(\Phi_1(u, v), \Phi_2(u, v)) - (\Phi_1(u', v'), \Phi_2(u', v'))\|_{\Sigma_\theta} \leq \frac{1}{2} \|(u, v) - (u', v')\|_{\Sigma_\theta}$$

Therefore the map $\Phi_1 \times \Phi_2 : \Sigma_\theta \rightarrow \Sigma_\theta$ is a contraction mapping, and we obtain a unique fixed point in Σ_θ which solves the equation (4.5).

Now we prove Theorem 1.1. Using the scaling (1.2) by taking

$$\lambda = \left(\frac{\epsilon_0}{2(\|v_0\|_{H^{-3/4}} + 1)} \right)^{4/3},$$

then we see

$$M = 2c_0 \|\phi_1\|_{L^2} = 2c_0 \lambda^{3/2} \|u_0\|_{L^2}, \quad \|\phi_2\| \leq \epsilon_0.$$

Thus we get local existence of the solution to (1.4) on $[0, T_1]$ for

$$T_1 \sim \left(\frac{\lambda^{3/2}}{M^2} \right)^{1/\theta} = \left(\frac{1}{\lambda^{3/2} \|u_0\|_{L^2}^2} \right)^{1/\theta}.$$

Therefore, for the original system (1.1), we get the local existence of the solution on $[0, T]$ for

$$T = \lambda^3 T_1 \sim T(\|u_0\|_{L^2}, \|v_0\|_{H^{-3/4}}).$$

Therefore, we obtain existence. By standard arguments, we can prove the rest parts of Theorem 1.1.

REFERENCES

- [1] D. Bekiranov, T. Ogawa and G. Ponce, Weak solvability and well-posedness of a coupled Schrödinger-Korteweg de Vries equation for capillary-gravity wave interactions, *Proceedings of the AMS.*, 125 (10), (1997), 2907-2919.
- [2] D. Bekiranov, T. Ogawa and G. Ponce, Interaction equations for short and long dispersive waves, *J. Funct. Anal.*, 158, (1998), 357-388.
- [3] J. Bourgain, Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations I, II. *Geom. Funct. Anal.*, 3:107-156, 209-262, 1993.
- [4] M. Christ, J. Colliander, and T. Tao, Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations. *Amer. J. Math.* 125 (2003), no. 6, 1235-1293.
- [5] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, Sharp global well-posedness for KdV and modified KdV on $\mathbb{R}$ and $\mathbb{T}$. *J. Amer. Math. Soc.* 16 (2003), 705-749.
- [6] A. J. Corcho, F. Linares, Well-posedness for the Schrödinger-Korteweg-de Vries system. *Trans. Amer. Math. Soc.* 359 (2007), no. 9, 4089-4106.
- [7] J. Ginibre, Y. Tsutsumi, and G. Velo, On the Cauchy problem for the Zakharov system. *J. Funct. Anal.* 151 (1997), no. 2, 384-436.
- [8] B. Guo, Ch. Miao, Well-posedness of the Cauchy problem for the coupled system of the Schrödinger-KdV equations, *Acta Math. Sinica, Engl. Series*, 15 (1999), 215-224.
- [9] Z. Guo, Local Well-posedness for dispersion generalized Benjamin-Ono equations in Sobolev spaces. *preprint*, arXiv:0812.1825
- [10] Z. Guo, Global Well-posedness of Korteweg-de Vries equation in $H^{-3/4}(\mathbb{R})$. *J. Math. Pures Appl.* 91 (2009), 583-597.

- [11] Z. Guo, B. Wang, Global well posedness and inviscid limit for the Korteweg-de Vries-Burgers equation, *J. Differential Equations* 246 (2009) 3864C-3901.
- [12] A. D. Ionescu, C. E. Kenig, Global well-posedness of the Benjamin-Ono equation in low-regularity spaces, *J. Amer. Math. Soc.*, 20 (2007), no. 3, 753-798.
- [13] M. Keel, T. Tao, Endpoint Strichartz estimates, *Amer. J. of Math.*, **120** (1998), 360-413.
- [14] C. Kenig, G. Ponce and L. Vega, Oscillatory integrals and regularity of dispersive equations, *Indiana University Math. J.*, Vol. 40, 1991, pp. 33-69
- [15] C. Kenig, G. Ponce, and L. Vega, The Cauchy problem for the Korteweg-de Vries equation in Sobolev spaces of negative indices. *Duke Math. J.* 71 (1993), no. 1, 1–21.
- [16] C. Kenig, G. Ponce, L. Vega, A bilinear estimate with applications to the KdV equation. *J. Amer. Math. Soc.*, 9:573-603, 1996. MR 96k:35159.
- [17] C. Kenig, G. Ponce, and L. Vega, On the ill-posedness of some canonical dispersive equations. *Duke Math. J.*, 106(3):617-633, 2001. MR 2002c:35265
- [18] H. Pecher, Hartmut Rough solutions of a Schrödinger-Benjamin-Ono system. *Differential Integral Equations* 19 (2006), no. 5, 517–535.
- [19] T. Tao, Multilinear weighted convolution of L^2 functions and applications to nonlinear dispersive equations. *Amer. J. Math.*, 123(5):839-908, 2001.
- [20] D. Tataru, Local and global results for wave maps I, *Comm. Partial Differential Equations* **23** (1998), 1781-1793.
- [21] M. Tsutsumi, Well-posedness of the Cauchy problem for a coupled Schrödinger-KdV equation, *Math. Sciences Appl.*, 2 (1993), 513-528.

LMAM, SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA
E-mail address: zihuaguo, wangyuzhao@math.pku.edu.cn